



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

diseased bark or both in the movements of the birds over these surfaces. This conclusion is supported by the fact that the birds tested were not carrying ascospores; that we have no evidence that ascospores are washed down the trees during the winter and spring months;¹¹ also that following a rain period pycnosporos are to be found in abundance on the healthy bark below blight lesions.

F. D. HEALD

R. A. STUDHALTER

FOREST PATHOLOGY LABORATORY,

U. S. DEPARTMENT OF AGRICULTURE,

PHILADELPHIA, PA.

THE RELATION BETWEEN ABNORMAL PERMEABILITY
AND ABNORMAL DEVELOPMENT OF *FUNDULUS*
EGGS

IN a previous paper¹ the suggestion was made that certain abnormalities in *Fundulus* embryos are caused by increase in permeability since osmotic pressure is not the cause and so many *different* substances have the *same* effect. It was found that the normal egg in distilled water or a "balanced" salt solution is impermeable to salts (Appendix II.). The egg appeared to be impermeable to water also, since enormous osmotic changes have no effect on it. The egg was found to contain nearly three times as much ash as sea water. The greater part of the ash is insoluble, but some of it may have been rendered so by the ashing. However, the soluble ash (3.18 per cent.) is as great as the total salts (2.84-3.29 per cent.) in the local sea water. And yet the egg develops normally, with little or no change in volume, in distilled water or in sea water that is evaporated to one half its volume, suggesting impermeability to water. The fact that the eggs dry up when exposed to air may be taken to indicate an increase in permeability to water, due to drying of the superficial layer or plasma membrane.

¹¹ Heald, F. D., and Gardner, M. W., "Preliminary Note on the Relative Prevalence of Pycnosporos and Ascospores of the Chestnut Blight Fungus during the Winter," SCIENCE, N. S., 37: 916-917, 1913.

¹ McClendon, *Am. Jour. Physiol.*, 1912, XXIX., p. 290.

In the same paper some preliminary chemical studies of the permeability were described, and the view advanced that the egg is normally impermeable to Mg ions, but since Mg was found to diffuse out of the eggs in a pure NaCl solution, this solution may have increased the permeability to Mg (p. 296). Only one experiment to test the permeability to anions was described. MgSO₄ solution was used, with negative results. However, the MgSO₄ contained too large a trace of chloride to make it possible to detect a very small diffusion of chloride from the eggs.

During the present season I was able to obtain especially pure salts, and have observed diffusion of both anions and cations from the eggs in pure solutions of these. The monstrosities produced in unbalanced salt solutions have also been studied. The experiments support the following generalizations:

1. Any solution of one or more of the salts of sea water, which is sufficiently unbalanced by other salts, *i. e.*, has a certain excess of some one cation, produces a number of types of monstrosities in *Fundulus* eggs. The types of monsters produced by the excess of one cation (*e. g.*, Na) are the same as those produced by any other (*e. g.*, K, Ca or Mg). Thus a qualitatively *specific action of a salt or ion does not exist*.

2. *These unbalanced salt solutions cause an increase in the permeability of the egg to salts.* This conclusion is based on the following data: The eggs in distilled water or in van't Hoff's solution (made with nitrates) lose no salts or ions that can be detected, except the ions of carbonic acid. On the contrary, the eggs give out salts or their ions in a mixture of NaCl and KCl or in pure solutions of the following salts: NaCl or nitrates of Na, K, Ca or Mg *in concentrations that do not kill the eggs during the experiment.* If the eggs are killed a more rapid diffusion takes place. The methods used will be published elsewhere.

J. F. MCCLENDON

U. S. BUREAU OF FISHERIES,

WOODS HOLE, MASS.,

July 25, 1913